

Soaking Germinated Oil Palm Seeds in Amino Boost Transit Max® to Assess Shoot and Root Response

Dual Chelate Fertilizer Pty Ltd.
PO Box 963, 162 New Guinea Road Robinvale VIC 3549, Australia
Correspondence: Research and Development Division
Email: info@dualchelate.com

Oil Palm trees are the most popular edible vegetable oil crop in the world (Aba and Paul, 2012). Known for its high oil content and efficiency, palm oil a common ingredient in many foods and other goods such as soaps and cosmetics (Gesteiro et al., 2019). However, due to the increasing demand for palm oil, the industry is facing social and environmental issues surrounding the sustainability of palm oil production and are exploring routes to improve continuously improve the efficiency of Oil Palm trees to match global demand. One way to do this is through improvements in shoot development and increasing root exploration and development to provide young Oil Palm seedlings with a strong base to grow and promote the absorption and utilization of nutrients more efficiently. This trial was done to assess if soaking pre-germinated Oil Palm seeds in 2% and 20% solutions of Amino Boost Transit Max® will increase shoot and root development compared to a control. It was found that though soakings of Amino Boost Transit Max, there was noticeable increases in shoot height compared to the control and also increases in root length compared to the control. It was also noted that seeds soaked in Amino Boost Transit Max® produced more lateral root increasing the surface area of the root system.

Key words: Oil Palm, pre-germinated seeds, Amino Boost Transit Max®, plant height, root length, lateral roots

Introduction

Oil Palm (*Elaeis guineensis*) trees are grown for the harvesting of the edible vegetable oil known as palm oil (Gesteiro et al., 2019). The palm oil is extracted using 2 different methods which utilize different parts of the palm seeds. These 2 methods created 2 different types of oil known as palm oil and palm kernel oil. Palm oil is traditionally used for edible purposes such as cooking, frying and as a food ingredient. Palm kernel oil is generally used for non-edible purposes such as the formulation of soaps, cosmetics and detergents (Chin, 2019).

For many years Palm Oil has been the most popular vegetable oil as it is the highest yielding vegetable oil crop which makes it very efficient and very popular. Another advantage is it requires much less land compared to other vegetable oil crops in order to produce the same amount of oil. Together, these advantages make oil palm production the most attractive vegetable producing crop also making it the least expensive vegetable oil in the world.

Oil Palm plantations are most commonly seen in the tropical zones around the equator where there is ample rainfall and warm weather. This tropical weather makes it a prime location for high yielding Palm Oil production. Indonesia produces the most Palm Oil with approximately 33.4 million tonnes produced annually, followed by Malaysia and Thailand. Indonesia and Malaysia make up approximately 84% of the market share (Which countries grow and produce palm oil?, 2015). However, over recent decades, there has been a strong increase in the demand for edible vegetable oil and as a result, Palm Oil plantations have had to expand rapidly in order to reach global demands. This increase in demands has significantly contributed to economic growth and poverty reduction in these countries, however many social and environmental issues have arisen related to the sustainability of the crop such as land degradation and deforestation (Qaim, Sibhatu, Siregar and Grass, 2020). Research and development is currently underway to explore avenues to further increase the efficiency of the Palm Oil trees to make more oil, make cultivars which are more suitable to arid environments where space is plentiful and also improve growing traits.

One way the efficiency of Palm Oil production can be increased is by reducing seedling mortalities, increasing shoot development and improving root exploration and development. By making improvements to the initial stages of growth from the germination of the Palm Oil seed to the development of the eophyll (first palm frond), seedlings can be transplanted quicker into the field and also have a better capability to absorb nutrients and accelerate growth making the palms more efficient and possibly shortening the time to first fruit production.

Dual Chelate Fertilizer Pty Ltd has developed a premium quality plant bio stimulant to assist in increasing root growth, improving the translocation of nutrients and enhancing the establishment of young plants. Amino Boost Transit Max® contains: 10% Amino Acids, 6%, Kelp, 4% Fulvic Acids, 1.5% Biologically Active Organic Molecules (BAOM) and 1.4% Nitrogen (Amino Acid derived Nitrogen). Amino Boost Transit Mix® in previous studies has proven to increase the root structure of numerous horticultural crops such as table grapes and almonds and also increase yields through strategic applications.

In this study, Amino Boost Transit Max® will be applied to germinated Palm Oil seeds through soaking in an effort to compare the development of the initial root system and shoot growth. Two different concentrations of Amino Boost Transit Max® soaking solutions will be used to examine any responses to the bio stimulant and then compared to control germinated Palm Oil seeds. Shoot height, root length and plant structure will be analyzed.

Objectives

1. Assess shoot growth and root system developmental differences between each Amino Boost Transit Max® soaking solution concentration against the control.
2. Measure shoot length, root length and take images of young Oil Palm seedlings 1 month after initial soaking.
3. Visually compare the treatments to see any difference in appearance.
4. Determine if Amino Boost Transit Max® is a beneficial product to use to improve the initial growth of Palm Oil seedlings.

Site Selection and Trial Design

The trial will be conducted in the greenhouse located at Dual Chelate Fertilizer. This site was selected as Oil Palms needs warm, humid and wet conditions. These conditions can be simulated using an irrigation sprinkler system to ensure that the germinated seeds do not dry out. Each pre-germinated Palm Oil seed is planted in its own individual.

In total, there will be 2 different seed soaking solutions of Amino Boost Transit Max® with concentrations at 2% and 20%. In each treatment there is 12 germinated Palm Oil seeds which are soaked for 12 hours before planting. See figure 1 for a diagram of the trial set up.



Figure 1: Soaking of germinated Oil Palm seeds in various concentration of Amino Boost Transit Max®. Germinated seeds were soaked for 12 hours then planted.

Once the seeds had been soaked for 12 hours in their respective solutions, the seeds were then planted root first in soil and covered with vermiculite to prevent moisture loss. The soil was collected from the vineyard at Dual Chelate fertilizer and is free from any nutrient contaminations. See figure 2 for the pot layout for the three different treatments (2% Amino Boost, 20% Amino Boost and control).




Figure 2: Pot set up of germinated Oil Palm seeds. each treatment has 12 pots with a single seed planted in each pot.

Once the Oil Palm seeds were planted, they were kept moist every day with short and frequent sprinkling. After 1 month, each plant was measured for its shoot height. The first plant from each treatment was also carefully dug up to examine the root system and measure the root length. One plant was examined as this experiment is on-going and will be again measured after 2 months.

Application Rates

Table 1: Application rates of different soaking solutions at 2% and 20% Amino Boost Transit Max(R). 12 Germinated Oil Palm seeds are soaked in each treatment for 12 hours then planted.

Treatment	Amino Boost Transit Max®	Water
2% Amino Boost	4ml	200ml
20% Amino Boost	40ml	200ml
Control	0ml	200ml

Observations

Shoot Height

Each plant was measured for its shoot height using a ruler after 1 month of growth. The shoot height is measured from the start of the root to the tip of the first palm frond (eophyll). Oil Palm seeds which did not grow was discounted from the measurements and not used when averaging the results for each treatment.

Root Length

The first plant from each treatment was then uprooted carefully and dirt was washed off. The longest root was then measured and photographed.

Comparative Photos

Comparative photos were taken to visually compare plant growth of the plants which were dug up to assess root quality and root length.

Results




Figure 3: Average plant height of germinated Oil Palm seeds soaked in Amino Boost 2% and Amino Boost 20% solution. Plant height was taken 1 month after planting.

Table 2: Root length of Oil Palm seedlings (cm)

Treatment	Root Length of 1 Plant (cm)
Control	8.1 cm
Amino Boost Transit Max® 2%	8.9 cm
Amino Boost Transit Max® 20%	10.1 cm



Figure 4: Image taken of germinated Oil Palm seedling soaked in a **2% Amino Boost Transit Max[®]** solution before planting for 12 hours compared to a control seedling. Photo taken one month after planting.



Figure 5: Image taken of germinated Oil Palm seedling soaked in a **20% Amino Boost Transit Max[®]** solution before planting for 12 hours compared to a control seedling. Photo taken one month after planting.

Discussion

When looking at figure 3, it can be seen that both the 2% and 20% Amino Boost Transit Max® treated Oil Palm seeds had taller shoots compared to the control. Although there was not much of a difference between the control and the 2% solution, there was a large increase in shoot height from 3.5cm to 4.7cm tall between the 2% and 20% solution. This is a percentage increase of 34%. Although there is no statistical significance between the control and the Amino Boost Transit Max® treated Oil Palm seedling heights, there is still a noticeable difference.

Amino Boost Transit Max® contains 10% organically derived amino acids made from 16 different types of amino acids, 6% Kelp, 4% Fulvic acid, 1.5% Biologically Active Organic Molecules (BAOM) and 1.4% amino acid derived nitrogen. When soaking the germinated Oil Palm seeds for 12 hours in the 2% and 20% Amino Boost solution these beneficial molecules and elements are able to be absorbed. Absorption is also enhanced due to the Biologically Activate Organic Molecules which facilitates in the movement via organic chelation of nutrients and molecules into cell which are then used in a number of different processes in the plant which can assist with the development of new growth. For example, Auxin is a growth promoting hormones which is produced in plants which orchestrates cell division, elongation and differentiation, apical dominance and also stimulates stem tropisms (Carey and Krogan, 2017). Auxin is made using amino acids such as tryptophan. Tryptophan is an amino acid which can be synthesized by the use of Amino Boost Transit Max®, therefore the Oil Palm seeds are able to convert this into Auxin which explains the increase in plant shoot height. Another benefit of providing amino acids is that the plants don't need to expel energy to make them and instead use energy reserves on more important aspects of plant growth.

Amino Boost Transit Max® also contains cytokinin based Kelp which is very beneficial in the early growth of plants as it contains a number of key elements such as Potassium and also high in alginic acid (Ali, Ramsabhag and Jayaraman, 2021). Since the kelp is already naturally high in Cytokinin (another key plant growth hormone), it is absorbed in the soaking process and subsequently used to promote cell division and also promotes apical dominance (Feng, Shi, Yang and Zuo, 2017). These benefits directly relate to plant height increases seen in figure 3.

Table 2 shows the root length of Oil Palm seeds soaked in Amino Boost Transit Max® compared to the control. It can be seen that there is a similar trend seen in the plant height results where there is not much of a difference seen between the control and 2% solution, however there is a large difference seen in the 20% solution. When looking at figure 5, it can actually be seen that the 20% treated Oil Palm seed root actually reaches the bottom of the pot and grows sideways. In previous research, it has shown that Amino Boost Transit Max® significantly increases the number of root hairs, lateral growth of roots and increases the endodermis diameter of the roots in establishing crops. This is because of the cytokinin based kelp has a direct effect on root apical dominance. This increase in apical dominance drives the main root deeper into the ground producing a longer root which is seen in both 2% solution and 20% solution.

Figure 4 and 5 show images of 2% and 20% Amino Boost treated seedlings compared to a control seedling respectively. These images show the increased shoot growth and also longer root development in the Amino Boost Transit Max® treated Oil Palm seeds. Additionally, there is more lateral root development in both concentrations seen on the main stem. These lateral roots increase the plants root system and increase the surface area for more nutrient and water

uptake. This is most likely a response from the Cytokinin in the Amino Boost Transit Max®. Also, by supplying the Oil Palm seeds with available amino acids, more energy was spent on cell elongation and division instead of using energy to synthesis large amino acid molecules.

Conclusion

In conclusion, this trial was conducted to assess the shoot and root response to soaking pre-germinated Oil Palm seeds in 2% and 20% solutions of Amino Boost Transit Max® for 12 hours. Both treatments showed that there was a positive response to the soaking of the seeds 1 month later by increasing plant height and also increasing root length whilst also improving lateral root production from the primary root.

When studying the figures and tables presented above, it can be found that Oil Palm seeds soaked in 2% and 20% Amino Boost Transit Max® solution for 12 hours had the following improvements compared to Oil Palm seeds soaked in water:

- Increased the plant height from 3.1cm (control) to 3.5cm (2% solution) and 4.7cm (20% solution)
- Increased root length from 8.1cm (control) to 8.9cm (2% solution) and 10.1cm (20% solution)
- Improved the root length and promoted the formation of lateral root development on seeds soaked in Amino Boost Transit Max® compared to the control.

This trial is still under observation with a second application of 2% and 20% Amino Boost Transit Max® solution being applied to the Oil Palm seedlings after 1 month of growth. After 2 months, shoot height and another analyses of root growth will be done to assess the further benefits of using Amino Boost Transit Max® to help increase establishment of Oil Palm seeds.

References

- Aba, Simon & Baiyeri, Paul. (2012). The oil palm.
- Ali, O., Ramsubhag, A. and Jayaraman, J., 2021. Biostimulant Properties of Seaweed Extracts in Plants: Implications towards Sustainable Crop Production. *Plants*, 10(3), p.531.
- Carey, N. and Krogan, N., 2017. The role of AUXIN RESPONSE FACTORS in the development and differential growth of inflorescence stems. *Plant Signaling & Behavior*, 12(4), p.e1307492.
- Chin, S., 2019. *Palm oil vs palm kernel oil: What's the difference?*. [online] Malaysiakini. Available at: <<https://www.malaysiakini.com/knownmypalmoil/480694>>
- Feng, J., Shi, Y., Yang, S. and Zuo, J., 2017. Cytokinins. *Hormone Metabolism and Signaling in Plants*, pp.77-106.
- Gesteiro, E., Guijarro, L., Sánchez-Muniz, F., Vidal-Carou, M., Troncoso, A., Venanci, L., Jimeno, V., Quilez, J., Anadón, A. and González-Gross, M., 2019. Palm Oil on the Edge. *Nutrients*, 11(9), p.2008.
- Green Palm. 2015. *Which countries grow and produce palm oil?*. [online] Available at: <<https://greenpalm.org/about-palm-oil/where-is-palm-oil-grown-2>>

Appendix 1




Figure 6: The plant height measurements taken from all treatments in the experiment. Red stars indicate statistically significant treatments